

Oxidative stress markers were measured in whole body homogenates of flies (not specifically in the brain or heads like of

changed to not relevant in distiller due to reason 3 081417
changed to foreign-language in distiller 081417
changed to foreign-language in distiller 081417
changed to not relevant in distiller due to reason 3 081417
removed animal tags and added in extra tag in distiller 081417
added animal primary tag in Distiller 081417
changed to not relevant in distiller due to reason 3 081417
added animal primary tag in Distiller 081417
changed to foreign-language in distiller 081417
ther Drosophila studies)
added animal primary tag in Distiller 081417

6230
6311
6326
6426
6505
6552
6579

J. Shajan A. Antony D. Cynthia Pauline P. Karthick Raja Namasivayam S. Nellore. Proteinaceous compounds from fragari
T. L. Yong V. W. Wall R. A. Jones K. Perry. PARAQUAT AND 2 ENDOGENOUS ANALOGS OF THE NEUROTOXIC SUBSTANCE
L. C. D. Carney C. Shen B. Wong S. Halaszynski K. Salomon M. P. Davies K. J. A. Tower J. Pomatto. The Mitochondrial Lon
J. Sirapat W. Thiantanawat A. Satayavivad. Neurological effects of chronic exposure to low doses of paraquat in rats. Rese
S. K. Dimri U. Kataria M. Kumari P. Singh. Ameliorative activity of withania somnifera root extract on paraquat-induced o
S. P. Salam S. K. N. Jaafar H. Gan S. H. Muzaimi M. Sulaiman S. A. Tang. Tualang Honey Protects the Rat Midbrain and Lun
M. Richfield E. K. Goodman B. M. Cory-Slechta D. A. Thiruchelvam. Developmental exposure to paraquat and maneb: effe